

Historic, archived document

Do not assume content reflects current scientific knowledge, policies, or practices.



76
Circular No. 643

November 1942 • Washington, D. C.

UNITED STATES DEPARTMENT OF AGRICULTURE

LIBRARY
RECEIVED
DEC 23 1942
U. S. Department of Agriculture

Owlet Moths (Phalaenidae) Taken at Light Traps in Kansas and Nebraska¹

H. H. WALKDEN, *assistant entomologist, Division of Cereal and Forage Insect Investigations, Bureau of Entomology and Plant Quarantine*, and D. B. WHELAN, *assistant entomologist, Nebraska Agricultural Experiment Station*

United States Department of Agriculture, Bureau of Entomology and Plant Quarantine, in Cooperation with the Nebraska Agricultural Experiment Station.

CONTENTS

	Page		Page
Introduction.....	1	Seasonal flight records.....	14
Traps.....	2	Flight records of individual species.....	18
Collections.....	3	Summary.....	18
Proportion of the sexes.....	14		

INTRODUCTION

The information contained in this publication was obtained through the operation of light traps located at six widely separated points in the Missouri Basin, in areas typical of both the semiarid and humid phases of the agriculture of this region. It deals with the seasonal occurrence and abundance of the owlet moths (Phalaenidae (Noctuidae)), a group composed largely of such notoriously injurious pests as the corn earworm, the pale western cutworm, the western armyworm, the wheat-head armyworm, and many other seriously injurious cutworms inhabiting the region surveyed. This region is of exceptional interest to workers in applied entomology because it contains an owlet moth fauna representing not only the strictly subterranean species, controllable only by modification of cultural practices, but those possessing either intermediate or surface-inhabiting characteristics, and more or less amenable to insecticidal treatment.

The data herein contained, revealing the distribution, seasonal flight periods, and peaks of abundance of the various species, are basic to, or of value in, the consideration of cultural control and other methods for the suppression of the various pests involved.

The use of light traps offered an efficient means of obtaining this information. Accordingly, in the fall of 1934 traps were set up in

¹ Published as Paper No. 223, Journal Series, Nebraska Agricultural Experiment Station;
433064°—42—1

several locations and their operation was continued during the flight seasons of 1935-37. A record of the nightly catch was kept, the species, number of moths captured, and the proportion of the sexes being tabulated. The traps were put in operation early enough in the spring so that the first flights could be observed, and they were kept in operation until cold weather in the fall stopped all flight activity.

Observations were made in the following six localities: Cherryvale, Garden City, Hays, and Manhattan in Kansas and Lincoln and Scottsbluff in Nebraska (fig. 1). These localities ranged in elevation




FIGURE 1.—Location of the light traps.

as follows: Cherryvale 800 feet, Garden City 2,800 feet, Hays 2,000 feet, Mannattan 1,100 feet, Lincoln 1,200 feet, and Scottsbluff 3,900 feet.

TRAPS

With the exception of the one used at Manhattan, the traps were of similar type and were designed by the senior writer. They consisted of an inverted galvanized-iron cone, 2 feet in diameter, with a roof of the same material. The lamp (fig. 2, *B*) was suspended in the center just above the rim, and a fruit jar was attached to the bottom of the cone for receiving the catch. The killing agent, calcium cyanide, was put in a small sack and placed in the fruit jar. The cyanide was renewed daily.

The trap used at Manhattan (fig. 2, *A*) was designed by R. C. Smith, of the Kansas Agricultural Experiment Station. Except for a more shallow cone, and the use of glass buffers about the lamp, this trap was the same as those at the other localities. Five-hundred-watt clear lamps were used at Manhattan and Lincoln and 200-watt inside-frosted lamps elsewhere.

The locations of the traps were probably not ideal, but the availability of electric outlets was the deciding factor in the selection of sites. These were as follows:

(1) At Cherryvale the location was unsatisfactory. It was on a flour mill that was illuminated nearly every night and near a transformer station that was always brilliantly lighted. The competition of these other lights greatly affected the catches, and the results from this trap are not considered of much value.

(2) At Garden City the trap was located on the grounds of the Garden City Branch Station, Kansas Agricultural Experiment Station, about 5 miles northeast of the city. The light was visible in all directions, although a small grove of trees on the southwest somewhat obscured it in that direction. It was placed about 5 feet above the ground.

(3) At Hays the trap was placed in an open space between buildings on the grounds of the Fort Hays Branch Station, Kansas Agricultural Experiment Station. It was about 5 feet above the ground and was visible for a long distance to the east and south.

(4) At Manhattan the trap was placed on top of an outbuilding near the college power plant. The lamp was about 20 feet above the ground and could be seen for




FIGURE 2.—Light trap used at Manhattan, Kans. (A), and at the other locations (B).

some distance to the west, south, and north, but was obscured by buildings on the east.

(5) At Lincoln the trap was placed at the east edge of the Nebraska Agricultural Experiment Station orchard, well within the city limits. It commanded a view to the east for some distance, but in other directions the light was obscured by the orchard trees nearby.

(6) At Scottsbluff the trap was located on the grounds of the Scottsbluff Branch Station, Nebraska Agricultural Experiment Station, about 6 miles northwest of the city. The trap was elevated about 10 feet above the ground and was visible for a long distance in all directions.

COLLECTIONS

The several light traps were in operation for various periods during the 4 years. Except for certain unavoidable interruptions, they were operated each night throughout the flight season.² The catches from Cherryvale, Garden City, Hays, and Scottsbluff were forwarded to

² The traps at Manhattan and Lincoln were operated by the writers and those at Cherryvale, Garden City, Hays, and Scottsbluff by the N. Sauer Milling Co., F. A. Wagner, L. C. Aicher, and John Carter, respectively.

either Manhattan or Lincoln, where the material was sorted and the results were tabulated. The number of nights for which records were obtained are as follows: Cherryvale 202, Garden City 612, Hays 214, Manhattan 629, Lincoln 656, and Scottsbluff 406.

It was at first intended to record all species of Phalaenidae taken at the several traps. Limitations of time, however, made it necessary to omit some of the smaller forms and the species of the genus *Catocala*, and in 1937 only species of known economic importance were recorded from Manhattan and Garden City. Altogether 305 species were taken and 525,547 individuals. The total yearly catch of each species for the different localities, together with the flight period, is shown in table 1. The sequence of genera and species is in accordance with McDunnough's Check list of Lepidoptera.³ Much of the material was identified by the writers, largely by comparison with specimens determined by J. F. Gates Clarke, of the Division of Insect Identification, Bureau of Entomology and Plant Quarantine, Rowland R. McElvare, New York City, and the late F. H. Benjamin. Credit is due these men for their careful and painstaking work.

³ McDUNNOUGH, J. CHECK LIST OF THE LEPIDOPTERA OF CANADA AND THE UNITED STATES OF AMERICA. PART I. MACROLEPIDOPTERA. South. Calif. Acad. Sci. Mem. 1,275 pp. 1938.

TABLE 1.—*Phalaenid moths caught in light traps at various localities in Kansas and Nebraska, 1934-37*—Continued

Species	Flight period	Number of moths caught													
		Cherryvale, Kans.		Manhattan, Kans.		Hays, Kans.		Garden City, Kans.		Lincoln, Nebr.		Scottsbluff, Nebr.			
		Mar. 8 to Oct. 31, 1935	Apr. 8 to Oct. 15, 1936	Sept. 26 to Oct. 30, 1934	Apr. 5 to Oct. 30, 1935	Sept. 28 to Nov. 23, 1934	Mar. 24 to Nov. 20, 1935	Apr. 10 to Oct. 31, 1936	Apr. 12 to Nov. 12, 1937	Mar. 14 to Oct. 28, 1935	Apr. 9 to Oct. 31, 1936	Apr. 13 to Oct. 30, 1937	July 13 to Oct. 19, 1935	May 3 to Oct. 31, 1936	May 17 to Oct. 18, 1937
Phalaeninae (Agrotinae)—Cont.															
<i>E. immitis</i> (Grt.)	May-July	227	312	774				1		25	132	152	1	8	21
<i>E. redipennis</i> (Grt.)	Sept.-Oct.	1								1	3				1
<i>E. scandens</i> (Riley)	May-Aug.										1	1	51	251	242
<i>E. mediata</i> (Grt.)	Oct.	3													
<i>E. placida</i> B. and McD	do														
<i>E. mimallonia</i> (Grt.)	Aug.	8	9	22	1	6	6	6	29	79	157	230	1	77	10
<i>E. messoria</i> (Harr.)	Sept.-Oct.				35	75	14	1	7				3	2	
<i>E. rectifincta</i> (Sm.)	Sept.-Oct.												14	342	80
<i>E. tessellata</i> (Harr.)	May-Aug.		1					1							
<i>E. tessellata tesselloides</i> (Grt.)	Oct.														
<i>E. remota</i> (Sm.)	July										3	1		37	
<i>E. simona</i> McD	June													1	
<i>E. declarata</i> (Walk.)	July-Sept.												30		
<i>E. verficilis</i> (Grt.)	July													1	
<i>E. albipennis</i> (Grt.)	June-Sept.														18
<i>E. obeliscoides</i> (Guen.)	July-Oct.	1				3				4	1	6			1
<i>E. costata thalocnisis</i> (Grt.)	July														
<i>E. clausa</i> McD	June														
<i>E. tristicula</i> (Morr.)	May-Aug.													2	
<i>E. tristicula nesitens</i> Sm	May							4							76
<i>Chorizagrotis auxiliaris</i> Grt.	Apr.-Oct.	1	307	1,769	560	2,839	18,062	28,300	12,775	90	210	288	285	909	1,020
<i>C. auxiliaris montana</i> Cook	July						16						1		
<i>C. thanatologia</i> (Dyar)	May-Oct.						1								
<i>C. thanatologia perfida</i> (Dodr.)	do						3								
<i>Protezarria balamitis</i> (Grt.)	do														
<i>Lozagrotis apicalis</i> (Grt.)	Sept.	20	14	12	31	31	1	1	1	5	1	3	2		
<i>L. albicosta</i> (Sm.)	Aug.-Oct.	3	139	96	101	1,221	98	25	31	39	250	1,111	169	661	402
<i>Onychagrotis rileyana</i> (Morr.)	Aug.-Oct.													1	1
<i>Eucrotopocnemis fimbriaris</i> (Guen.)	Oct.														
<i>E. tripars</i> (Walk.)	do														
<i>Agrotis vastula</i> Walk.	Sept.-Oct.	14	10	2	14	101	23	562	108	414	4	4	147	161	65
<i>A. areolata</i> (Sm.)	Apr.-June	2	1			3	3	56	49	3	1			39	14
<i>A. orthogonia</i> Morr.	Aug.-Oct.					36	14	395	24	69	489		16	64	19
<i>A. gladiaria</i> Morr.	Sept.-Oct.	34	185	310	208	4,062	2		69	77	316	291	4	4	2

	156	276	459	428	3,171	892	100	71	2	3	679	1,171	2,701	240	413	211
<i>A. tenerabilis</i> Walk.																
<i>A. volubilis</i> Harv.	1	2	2	3	2	2	26	34	3	73	13				126	38
<i>A. maleficata</i> Guen.	131	14	201	53	382	28	93	560	800	1,179	768	107	620	24	14	17
<i>A. ypsilon</i> (Rott.)	228	1,031	4,544	1,963	1,622	1,737	2,967	52	198	2,688	341	1,181	3,487	292	279	186
<i>Faitia subgibata</i> (Haw.)																
<i>F. annexa</i> (Treat.)	90	4,942	92	74	33		3	10	3	49	10	9	24	3	61	1
<i>Copablepharon grandis</i> Stkr.									3	5						
<i>C. contrastata</i> McD.																
<i>C. longipenne</i> Grt.																
<i>C. alba</i> (Harv.)																
<i>Spacotlis claudestina</i> (Harr.)																
<i>S. hawaiiæ</i> (Grt.)																
<i>Chœrophora fungorum</i> G. and B.	49	2	14					1								
<i>Anicla infacta</i> (Ochs.)			1	2		1							1			
<i>Hemiteuzia rudens</i> (Harv.)			1	2			3	6	9				11			4
and <i>i. pellucidata</i> (Grt.)																
<i>Cerastis tenaxifera</i> (Walk.)																
<i>Pteridroma margaritosa</i> Haw. and <i>i. saucia</i> (Hbn.)	93	22	761	356	4,683	6	77	49	727	1,055	455	420	2,197	77	71	357
<i>Caradrina quadrangula clemens</i> (Sm.)																
<i>Amalthea nigrum</i> (L.)	1	1	47	123	113			1	17	16	50	21	48	27	48	8
<i>A. badinoides</i> (Grt.)	39	24	41	26	41	2					12	30	25			
<i>Pseudoglossa olivata</i> (Harv.)																
<i>Abagrotis placida</i> (Grt.)																
<i>A. barnesi</i> (Bcni.)	1			3	2			3						5	8	10
<i>Rhynchagrotis cupida</i> (Grt.)																
<i>Ulex saturatus</i> Grt.								3								
<i>U. pictus</i> Grt.													1			
Hadennæ:																
<i>Scotogramma trifolii</i> (Rott.)	153	861	3,677	1,100	984	89	3,615	498	86,661	75,021	42,325	970	2,492	1,396	742	403
<i>S. trifolii albifusa</i> (Walk.)																
<i>Trichocla postica</i> Sm.																
<i>T. artesa</i> Sm.																
<i>Polia legitima</i> (Grt.)	4	6	15	5											10	
<i>Lacinipolia meditata</i> (Grt.)																
<i>L. anguina</i> (Grt.)	3	3	9	21			76				1	4	28	1	1	3
<i>L. incurva</i> (Sm.)	5										57	102				1
<i>L. vicina</i> (Grt.)																
<i>L. renigera</i> (Steph.)	33	32	43	7		233	101		2	1			2	4	7	2
<i>L. ereda</i> (Walk.)		2	23	65							211	1,011	60	7	5	3
<i>L. lora</i> (Guen.)																
<i>L. olivacea</i> (Morr.)																
<i>L. maritima</i> (Harv.)																
<i>Ancpia capsularis</i> (Guen.)	2	5	8	2												
<i>Trichotia signata</i> (Walk.)			1	2												
<i>Uloonche calca</i> (Guen.)			2													
<i>U. modesta disticha</i> (Morr.)																
<i>Orthodes akalus</i> (Stkr.)						2										
<i>O. incincta</i> (Morr.)						1										
<i>O. saturnus</i> (Stkr.)	16	51	104	83	141	10	16		1		36	91	103	117	98	9
<i>O. crenulata</i> (Bul.)	3			6								3			8	

	180	1,676	1,946	1,695	1,829	1,062	487	701	1,035	1,219	2,186	266	610	2,120	45	25	49
<i>H. armigera</i> (Hbn.)																	
<i>H. nitescens</i> (F.)			4	4			4	4	8				4	1	7	6	2
<i>Camphidia scutosa</i> (Schiff.)		11	6	8		1	2	4	9			1	2	2	1	7	1
<i>Oryzias citrinellus</i> (G. and R.)			7	1			1	58	2				3		1	68	2
<i>Chlorocleptria simplex</i> Sm.		1	5	10			7	158	167			1	20	21	5	54	
<i>Dasypodactea taecens</i> (Morr.)			30	67			4	140	195			1	3	1	11	209	
<i>D. needi</i> (Grt.)			3	11			1					4	3	3	5	3	
<i>Rhodophora gaurae</i> (A. and S.)	3		94	17			1					4	11	14	4	2	
<i>Thyreon rosea</i> Sm.				4									1	2	1	2	
<i>Schinia cumatilis</i> (Grt.)			131	17	13				6							3	4
<i>S. hulstia</i> Tepper				3												1	2
<i>S. reniformis</i> Sm.			3	33	5				1				9	1	1	2	4
<i>S. chryselia</i> (Grt.)			3	33	5				1				7	4	3	4	4
<i>S. trifascia</i> Hbn.					6								3	3	1	1	
<i>S. oleagina</i> Morr.												1	3	2	7	4	
<i>S. gracilenta</i> Hbn.			42	120			179	13	27								
<i>S. acutilinea</i> (Grt.)					3												3
<i>S. diffusa</i> Sm.																	
<i>S. walsinghami</i> (Hy. Edw.)																	
<i>S. nundina</i> (Dru.)			1														3
<i>S. ferta</i> (Grt.)	36		2,043	6,624			1,024	17	106			33	334	79	1	4	6
<i>S. albifascia</i> Sm.				1			3	2	50							4	4
<i>S. regia</i> (Stkr.)	3		5												1	4	4
<i>S. gloriosa</i> (Stkr.)					1											1	
<i>S. sanguinea</i> (Geyer)																	
<i>S. saturata</i> (Grt.)			13	268			3		13				2	2			
<i>S. thoreani</i> (G. and R.)	21		10	53	10												
<i>S. marginata</i> (Haw.)	30		8	4			1		2				2				
<i>S. tora</i> Stkr.				6										4			
<i>S. bifascia</i> Hbn.					5												
<i>S. ultima</i> (Stkr.)				10													
<i>S. jaguarina</i> (Guen.)	7		965	73			25	27	16			22	27	14	8	7	3
<i>S. ignis</i> (Guen.)													1	1			
<i>S. arcigera</i> (Guen.)	2	1	6	11			1									3	
<i>S. bicuspida</i> Sm.			13	36	2		1		2				1			3	2
<i>S. mortua</i> (Grt.)					3							2				3	3
<i>S. mortua packardii</i> (Grt.)	3		8	155			5	15	8						3	16	1
<i>S. do</i>		1															
<i>S. mortua nobilis</i> (Grt.)				27	22												
<i>S. sordida</i> Sm.			4	48													
<i>S. siren</i> (Stkr.)																	
<i>S. roseincta</i> (Harv.)				1													
<i>S. do</i>																	
<i>Trichostellus cupes</i> (Grt.)					1				6							4	1
<i>Erastria synochitis</i> G. and R.																	
<i>E. carnicola</i> Guen.			1	77						1					2		2
<i>Neorastria apicosa</i> (Haw.)				2										314			
<i>Chamyris cernitha</i> (Trelt.)																	
<i>Amyna octo</i> (Guen.)			1														
<i>Tarachidia erastrioides</i> (Guen.)																	
<i>T. tenuescens</i> (Sm.)																	
<i>T. semiflava</i> (Guen.)														944			10
<i>do</i>																22	1
<i>do</i>																	

Acontinae:

Although the response to light varies with different species, it is probable that the number of individuals recorded is a fairly reliable estimate of the relative abundance of the species. Approximately 90 percent of the total catch was referable to species of economic importance.

PROPORTION OF THE SEXES

During the course of this study an attempt was made to determine the proportion of the sexes in representative samples of the various species taken. In many cases, however, insufficient material was examined or captured for a reliable estimate to be obtained. The results are presented in table 2. Only those species represented by 25 or more specimens are tabulated, all others being included in one group at the end of the table. Altogether, 36,692 specimens were examined, and of this number 12,902, or 35 percent, were females. It will be seen that the proportion of females varied greatly with different species, ranging from 3 to 60 percent.

TABLE 2.—Proportion of females among various species of phalaenid moths caught

Species	Moths examined	Females	Species	Moths examined	Females
	Number	Percent		Number	Percent
<i>Acronicta lepusculina</i> Guen.	28	36	<i>Nephelodys emmedonia</i> (Cram.)	32	3
<i>A. parallela</i> (Grt.)	34	35	<i>Protoleucania albilinea</i> (Hbn.)	918	21
<i>Euxoa siccata</i> (Sm.)	256	10	<i>Leucania phragmatidicola</i>		
<i>E. niveilinea</i> (Grt.)	292	14	Guen.	170	46
<i>E. immixta</i> (Grt.)	50	22	<i>L. unipuncta</i> (Haw.)	4,160	36
<i>E. scandens</i> (Riley)	40	30	<i>Cucullia laetifica</i> Lint.	28	25
<i>E. messoria</i> (Harr.)	31	33	<i>Oncocnemis sanina</i> Sm.	79	19
<i>E. reticincta</i> (Sm.)	83	29	<i>O. augustus</i> Harv.	31	35
<i>E. tessellata</i> (Harr.)	32	44	<i>O. occata</i> (Grt.)	25	60
<i>Chorizagrotis auriliaris</i> (Grt.)	10,672	51	<i>Psaphidia grotei</i> (Morr.)	26	23
<i>Loragrotis apicalis</i> (Grt.)	32	31	<i>Crymodes burgessi</i> (Morr.)	502	6
<i>Onychagrotis rileyana</i> (Morr.)	262	13	<i>Platyperigea ertima</i> (Walk.)	30	7
<i>Agrotis vetusta</i> Walk.	293	28	<i>Prodenia ornithogalli</i> Guen.	1,501	25
<i>A. daedalus</i> (Sm.)	62	15	<i>Laphygma frugiperda</i> (A. and S.)	44	48
<i>A. orthogonia</i> Morr.	233	18	<i>L. erigua</i> (Hbn.)	510	32
<i>A. gladiaria</i> Morr.	185	27	<i>Selicanis cinereola</i> (Guen.)	67	27
<i>A. venerabilis</i> Walk.	984	3	<i>Heliethis paradoxa</i> (Grt.)	32	56
<i>A. malefida</i> Guen.	35	43	<i>H. armigera</i> (Hbn.)	3,149	45
<i>A. ypsilon</i> (Rott.)	1,012	32	<i>Schinia mortua packardii</i> (Grt.)	29	21
<i>Feltia subgothica</i> (Haw.)	2,767	7	<i>Autographa falcifera simplex</i>		
<i>F. annexa</i> (Treit.)	138	17	(Guen.)	59	8
<i>Peridroma margaritosa saucia</i>			<i>A. brassicae</i> (Riley)	528	39
(Hbn.)	936	36	<i>Caenurgina crassiuscula</i>		
<i>Amathes c-nigrum</i> (L.)	44	52	(Haw.) and <i>C. erecta</i>		
<i>A. hadinodis</i> (Grt.)	27	11	(Cram.) mixed	1,019	33
<i>Scotogramma trifolii</i> (Rott.)	3,550	35	All other species	1,133	33
<i>Laciniptolia meditata</i> (Grt.)	74	12			
<i>L. vicina</i> (Grt.)	272	13			
<i>L. renigera</i> (Steph.)	51	12			
<i>Orthodes incincta</i> (Morr.)	145	17			
			Total	36,692	35

SEASONAL FLIGHT RECORDS

Taken as a group, the light-trap records show that Phalaenidae are in flight from early in March to late in November, with periods of heavy flight and scarcity interspersed irregularly throughout the season. The nightly flight of Phalaenidae in the different localities is shown graphically in figures 3, 4, and 5. The nightly flight at Garden City was much greater than at any of the other localities, owing chiefly to the enormous numbers of adults of *Scotogramma trifolii* captured. The numbers of moths taken varied greatly in the different localities, and from year to year, and also as to the time of the year when the




FIGURE 3.—Nightly flight of Phalaenidae at Garden City, Kans., 1935-37.




FIGURE 4.—Nightly flight of Phalaenidæ at Manhattan, Kans., 1934-37.

period of greatest flight occurs. For example, at Garden City in 1935 the peak flight occurred in July, in 1936 in May, and in 1937 in June.




FIGURE 5.—Nightly flight of Phalaenidae at Lincoln and Scottsbluff, Nebr., and at Hays and Cherryvale, Kans., 1934-37.

In other words, there is no definite pattern in the flight periods of the Phalaenidae as a group, largely because individual species vary greatly in numbers from year to year.

FLIGHT RECORDS OF INDIVIDUAL SPECIES

The species of Phalaenidae fall into two main groups based on their seasonal history, namely, those species producing a single generation annually and those producing more than one generation annually.

As noted previously, species of economic importance comprised 90 percent of the total number of individuals captured. The seasonal flight of several single-generation species of economic importance is shown in figures 6 and 7.

In general, the flight of individual species showed the same trend at all the localities, the chief difference being in numbers taken. Where no striking differences were noted only one locality was plotted on the charts. It will be seen that the species producing a single generation annually have a rather short flight period, usually late in August, extending through September and into the early part of October. The notable exception is *Chorizagrotis auxiliaris*. The enormous flight of this species in May and June suddenly subsides and the species reappears again in September, after an estivation period of about 2 months. The fall flight is greatly reduced in numbers. This species is normally confined to the semiarid regions, but occasionally occurs in abundance at Manhattan (fig. 6, *B*) and at Lincoln. The records for the individual species show the same trend, namely, years of abundance usually followed by years of scarcity.

The seasonal flight of several species producing multiple generations annually is shown in figures 8, 9, 10, and 11. The flight records alone do not delineate clearly the various generations owing to overlapping as the season progresses. The charts, however, bring out some points of interest with reference to the seasonal abundance of multiple-generation species. *Heliothis armigera* (fig. 10, *A*) is shown to be scarce early in the season, with a marked increase in abundance in the fall. With *Prodenia ornithogalli* (fig. 10, *B*) and *Peridroma margaritosa saucia* (fig. 11, *A*) the peak of abundance is reached in midseason, followed by comparative scarcity in the fall. *Protoleucania albilinea* (fig. 11, *B*) shows two distinct flight periods, one early in May and the other late in August or early in September.

SUMMARY

Light traps were operated for various periods in six localities in Kansas and Nebraska during the period 1934-37, in order to obtain information on the flight periods and abundance of owlet moths (Phalaenidae). An attempt was made to operate the traps each night during the flight season, which extended from March to November. A total of 2,719 nightly records were made.

During the 4-year period 305 species of Phalaenidae were taken, totaling more than 525,000 individuals. Approximately 90 percent of the specimens taken were species of economic importance. The different species taken in each locality are shown, together with the total yearly flight, the flight period, and the proportion of the sexes. More than 36,000 individuals were examined for sex, and it was found that 35 percent were females. The sex ratio varied greatly with the species.

The seasonal flight records are presented graphically for the Phalaenidae as a group and for several species of economic importance. The numbers of moths taken differed greatly in the different localities,




FIGURE 6.—Seasonal flight of *Chorizagrotis auxiliaris* at Garden City, Kans. (A), and at Manhattan, Kans. (B), 1935-37.




FIGURE 7.—Seasonal flight of *Feltia subgothica* at Manhattan, Kans. (A), and at Lincoln, Nebr. (B), *Agrotis venerabilis* at Lincoln (C), *A. gladiaria* at Manhattan (D), *A. orthogonia* at Garden City, Kans. (E), and *Euxoa niveilinea* at Garden City (F), 1934-37.




FIGURE 8.—Seasonal flight of *Scotogramma trifolii* at Garden City, Kans., 1935-37.




FIGURE 9.—Seasonal flight of *Caenurgina erecta* and *C. crassiuscula* (A), and *Leucania unipuncta* (B), at Manhattan, Kans., 1935-37.




FIGURE 10.—Seasonal flight of *Heliothis armigera* at Manhattan, Kans. (A), and of *Prodenia ornithogalli* at Manhattan (B), and at Lincoln, Nebr. (C). 1935-37.




FIGURE 11.—Seasonal flight of *Peridroma margaritosa saucia* (A) and *Protoleucania albilinea* (B) at Manhattan, Kans., 1935-37.

and from year to year, and also as to the time of year when the greatest period of flight occurred. In general, however, flights of individual species showed the same trend in all the localities.

Species having a single generation annually had a rather short flight period, usually from late August to early October. *Chorizagrotis auxiliaris*, an exception to this rule, had an enormous flight in May and June and a much lighter flight in the fall.

Owing to overlapping of broods, the flights of the different generations of multiple-generation species were not clearly defined. *Heliothis armigera* was scarce early in the season, with a marked increase in abundance in the fall. Peak abundance of *Prodenia ornithogalli* and *Peridroma margaritosa saucia* occurred in midseason, followed by comparative scarcity in the fall. *Proteoleucania albilinea* showed two distinct flight periods, one early in May and the other late in August or early in September.

ORGANIZATION OF THE UNITED STATES DEPARTMENT OF AGRICULTURE WHEN THIS PUBLICATION WAS EITHER FIRST PRINTED OR LAST REVISED

<i>Secretary of Agriculture</i> -----	CLAUDE R. WICKARD.
<i>Under Secretary</i> -----	PAUL H. APPLEBY.
<i>Assistant Secretary</i> -----	GROVER B. HILL.
<i>Chief, Bureau of Agricultural Economics</i> -----	HOWARD R. TOLLEY.
<i>Director of Agricultural War Relations</i> -----	SAM BLEDSOE.
<i>Director of Extension Work</i> -----	M. L. WILSON.
<i>Director of Finance</i> -----	W. A. JUMP.
<i>Director of Foreign Agricultural Relations</i> -----	L. A. WHEELER.
<i>Director of Information</i> -----	MORSE SALISBURY.
<i>Director of Personnel</i> -----	T. ROY REID.
<i>Land Use Coordinator</i> -----	ERNEST H. WICKING.
<i>Librarian</i> -----	RALPH R. SHAW.
<i>Solicitor</i> -----	ROBERT H. SHIELDS.
<i>Chief, Office of Civilian Conservation Corps Activities</i> -----	FRED MORRELL.
<i>Chief, Office of Plant and Operations</i> -----	ARTHUR B. THATCHER.
<i>Administrator of Agricultural Marketing</i> -----	ROY F. HENDRICKSON.
<i>Administrator of Agricultural Conservation and Adjustment.</i>	M. CLIFFORD TOWNSEND.
<i>Chief, Agricultural Adjustment Agency</i> -----	FRED S. WALLACE.
<i>Chief, Soil Conservation Service</i> -----	HUGH H. BENNETT.
<i>Manager, Federal Crop Insurance Corporation</i> -----	LEROY K. SMITH.
<i>Chief, Sugar Division</i> -----	JOSHUA BERNHARDT.
<i>Administrator of Agricultural Research</i> -----	E. C. AUCHTER.
<i>Chief, Bureau of Animal Industry</i> -----	JOHN R. MOHLER.
<i>Chief, Bureau of Agricultural Chemistry and Engineering.</i>	W. W. SKINNER, <i>Acting.</i>
<i>Chief, Bureau of Dairy Industry</i> -----	O. E. REED.
<i>Chief, Bureau of Entomology and Plant Quarantine.</i>	P. N. ANNAND.
<i>Chief, Office of Experiment Stations</i> -----	JAMES T. JARDINE.
<i>Chief, Bureau of Plant Industry</i> -----	R. M. SALTER.
<i>Chief, Bureau of Home Economics</i> -----	LOUISE STANLEY.
<i>Superintendent, Beltsville Research Center</i> -----	C. A. LOGAN.
<i>President, Commodity Credit Corporation</i> -----	J. B. HUTSON.
<i>Administrator of Farm Security Administration</i> -----	C. B. BALDWIN.
<i>Governor of Farm Credit Administration</i> -----	ALBERT G. BLACK.
<i>Chief, Forest Service</i> -----	EARLE H. CLAPP, <i>Acting.</i>
<i>Administrator, Rural Electrification Administration</i> -----	HARRY SLATTERY.

This bulletin is a contribution from

<i>Bureau of Entomology and Plant Quarantine</i> -----	P. N. ANNAND, <i>Chief.</i>
<i>Division of Cereal and Forage Insect Investigations.</i>	C. M. PACKARD, <i>Principal Entomologist, in Charge.</i>



[